



Hemophilia Products – Factor VIII:

Advate, Adynovate, Afstyl, Eloctate, Hemofil M, Koate/Koate DVI, Kogenate FS, Kovaltry, Novoeight, Nuwiq, Obizur, Recombinate, Xyntha/Xyntha Solofuse, Jivi, Esperoct, Altuviio (Intravenous)

Document Number: IC-0340

Last Review Date: 06/01/2023

Date of Origin: 12/16/2014

Dates Reviewed: 12/2014, 04/2015, 05/2015, 09/2015, 12/2015, 03/2016, 06/2016, 12/2016, 06/2017, 09/2017, 11/2017, 09/2018, 10/2018, 03/2019, 10/2019, 02/2020, 09/2020, 06/2022, 04/2023, 06/2023

I. Length of Authorization

Coverage is provided for 3 months and may be renewed thereafter, unless otherwise specified*.

Note: The cumulative amount of medication the patient has on-hand will be taken into account for authorizations. Up to 5 'on-hand' doses for the treatment of acute bleeding episodes will be permitted at the time of the authorization request.

** Initial and renewal authorization periods may vary by specific covered indication*

II. Dosing Limits

A. Quantity Limit (max daily dose) [NDC unit]:

- N/A

B. Max Units (per dose and over time) [HCPCS Unit]:

- Advate: 73,600 billable units per 28-day supply
- Adynovate: 36,800 billable units per 28-day supply
- Afstyl: 69,000 billable units per 28-day supply
- Eloctate: 40,250 billable units per 30-day supply
- Kogenate: 43,125 billable units per 30-day supply
- Kovaltry: 86,250 billable units per 30-day supply
- Novoeight: 82,800 billable units per 28-day supply
- Nuwiq: 86,250 billable units per 30-day supply
- Hemofil M: 55,200 billable units per 28-day supply
- Koate DVI: 55,200 billable units per 28-day supply
- Recombinate: 55,200 billable units per 28-day supply
- Xyntha/Xyntha Solofuse: 41,400 billable units per 28-day supply
- Obizur: 115,000 billable units per 90-day supply

- Jivi: 41,400 billable units per 30-day supply
- Esperoct: 40,250 billable units per 28 days
- Altuviiiio: 23,000 units per 28 days

III. Initial Approval Criteria ^{1-17,22,23}

Hemophilia Management Program
Requirements for half-life study and inhibitor tests are a part of the hemophilia management program. This information is not meant to replace clinical decision making when initiating or modifying medication therapy and should only be used as a guide.

Coverage is provided in the following conditions:

A. Advate, Eloctate **Φ**, Hemofil M, Koate/Koate DVI, Kogenate FS **Φ**, Novoeight, Recombinate, Xyntha/Xyntha Solofuse **Φ**, Nuwiiq, Adynovate, Kovaltry, Afstyla, Jivi, Esperoct, Altuviiiio

Hemophilia A (congenital factor VIII deficiency) †

- Diagnosis of congenital factor VIII deficiency has been confirmed by blood coagulation testing; **AND**
- If the request is for Jivi, patient must be at least 12 years of age, or if request is for Altuviiiio, patient must be at least 1 year of age; **AND**
- Will not be used for the treatment of von Willebrand's disease; **AND**
- Used as treatment in at least one of the following:
 - On-demand treatment and control of bleeding episodes **OR**
 - Perioperative management (**Authorizations valid for 1 month*); **OR**
 - Routine prophylaxis; **AND**
 - Used to reduce the frequency of bleeding episodes; **OR**
 - Used to reduce the frequency of bleeding episodes and reduce the risk of joint damage in children without pre-existing joint damage (**Kogenate-FS ONLY**); **AND**
 - Patient must have severe hemophilia A (factor VIII level of <1%); **OR**
 - Patient has at least two documented episodes of spontaneous bleeding into joints.

Hemophilia Management Program
• If the request is for routine prophylaxis and the requested dose exceeds dosing limits under part II or if member BMI ≥ 30, a half-life study should be performed to determine the appropriate dose and dosing interval. • If the request is for Eloctate, Adynovate, Jivi, Esperoct, or Altuviiiio the following criteria should be met: ○ Patient is not a suitable candidate for a standard non- EHL factor VIII product. ○ A half-life study must be scheduled to determine the appropriate dose and dosing interval of the EHL product when initiated.

FACTOR VIII_HEMOPHILIA PRODUCTS - Prior Auth Criteria

Proprietary Information. Restricted Access – Do not disseminate or copy without approval.

©2023, Magellan Rx Management

- Prior to switching to Eloctate, Adynovate, Jivi, or Esperoct a half-life study should also be performed on current non- EHL factor VIII product to ensure that a clinical benefit will be achieved.
- If the request exceeds any of the following dosing limits, documentation must be submitted specifying why the member is not a suitable candidate for Hemlibra and alternative EHL factor VIII products.
 - 50 IU/kg every 4 days (total weekly dose of 87.5 IU/kg) for Eloctate
 - 40 IU/kg twice weekly (total weekly dose of 80 IU/kg) for Adynovate
 - 60 IU/kg every 5 days (total weekly dose of 84 IU/kg) for Jivi
 - 50 IU/kg every 4 days (total weekly dose of 87.5 IU/kg) for Esperoct
- For minimally treated patients (< 50 exposure days to factor products) previously receiving a different factor product, inhibitor testing is required at baseline, then at every comprehensive care visit (yearly for the mild and moderate patients, semi-annually for the severe patients)

B. Obizur¹⁰

Acquired Hemophilia A (acquired factor VIII deficiency) † Φ

- Patient is at least 18 years of age; **AND**
- Diagnosis of acquired factor VIII deficiency has been confirmed by blood coagulation testing; **AND**
- Used as on-demand treatment and control of bleeding episodes; **AND**
- Is NOT being used for congenital Hemophilia A OR von Willebrand disease; **AND**
- Patient does not have baseline anti-porcine factor VIII inhibitor titer >20 Bethesda Units (BU)

Hemophilia Management Program
● For members with a BMI ≥ 30, a half-life study should be performed to determine the appropriate dose and dosing interval. ● For minimally treated patients (< 50 exposure days to factor products) previously receiving a different factor product, inhibitor testing is required at baseline, then at every comprehensive care visit (yearly for the mild and moderate patients, semi-annually for the severe patients)

† FDA Approved Indication(s); ‡ Compendia Recommended Indication(s); Φ Orphan Drug

IV. Dispensing Requirements for Rendering Providers (Hemophilia Management Program)

- Prescriptions cannot be filled without an expressed need from the patient, caregiver, or prescribing practitioner. Auto-filling is not allowed.
- Monthly, rendering provider must submit for authorization of dispensing quantity before delivering factor product. Information submitted must include:
 - Original prescription information, requested amount to be dispensed, vial sizes available to be ordered from the manufacturer, and patient clinical history (including patient product inventory and bleed history)
 - Factor dose should not exceed +1% of the prescribed dose and a maximum of three vials may be dispensed per dose. If unable to provide factor dosing within the required

threshold, below the required threshold, the lowest possible dose able to be achieved above +1% should be dispensed. Prescribed dose should not be increased to meet assay management requirements.

- The cumulative amount of medication(s) the patient has on-hand should be taken into account when dispensing factor product. Patients should not have more than 5 extra doses on-hand for the treatment of acute bleeding episodes.
- Dispensing requirements for renderings providers are a part of the hemophilia management program. This information is not meant to replace clinical decision making when initiating or modifying medication therapy and should only be used as a guide.

V. **Renewal Criteria** ^{1-17,22,23}

Coverage can be renewed based upon the following criteria:

- Patient continues to meet indication-specific relevant criteria identified in section III; **AND**
- Absence of unacceptable toxicity from the drug. Examples of unacceptable toxicity include: anaphylaxis and hypersensitivity reactions (e.g., angioedema, chest tightness, dyspnea, wheezing, urticaria, pruritus, hypotension, etc.), thromboembolic events (thromboembolism, pulmonary embolism), development of neutralizing antibodies (inhibitors), etc.; **AND**
- Any increases in dose must be supported by an acceptable clinical rationale (i.e., weight gain, half-life study results, increase in breakthrough bleeding when patient is fully adherent to therapy, etc.); **AND**
- The cumulative amount of medication(s) the patient has on-hand will be taken into account when authorizing. The authorization will allow up to 5 doses on-hand for the treatment of acute bleeding episodes as needed for the duration of the authorization; **AND**

On-demand treatment of bleeding episodes and control of bleeding episodes

- Renewals will be approved for a 6-month authorization period.

Perioperative management of bleeding

- Coverage may NOT be renewed

Routine prophylaxis

- Renewals will be approved for a 12-month authorization period; **AND**
- Patient has demonstrated a beneficial response to therapy (i.e., the frequency of bleeding episodes has decreased from pre-treatment baseline)

VI. **Dosage/Administration** ^{1-16,22}

Advate

Indication	Dose
On-demand treatment and control of bleeding episodes Congenital Hemophilia A	Dose (IU/kg) = desired factor VIII rise (IU/dL or % of normal) x 0.5 (IU/kg per IU/dL) <u>Minor</u> Circulating Factor VIII required (% of normal) (20-40%) = 10-20 IU/ kg -Repeat every 12-24 hours as needed (every 8 to 24 hours for patients underage of 6). Continue until the bleeding episode is resolved (as indicated by relief of pain) or healing is achieved (approximately 1 to 3 days). <u>Moderate</u> Circulating Factor VIII required (% of normal) (30-60%) = 15-30 IU/ kg - Repeat every 12-24 hours as needed (every 8 to 24 hours for patients underage of 6). Continue until the bleeding episode is resolved (as indicated by relief of pain) or healing is achieved (approximately 3 days or more). <u>Major</u> Circulating Factor VIII required (% of normal) (60-100%) = 30-50 IU/ kg - Repeat every 8-24 hours as needed (every 6 to 12 hours for patients underage of 6). Continue until the bleeding episode is resolved.
Routine prophylaxis Congenital Hemophilia A	For prophylaxis regimen to prevent or reduce frequency of bleeding episodes, dose between 20 to 40 IU per kg every other day (3 to 4 times weekly). Alternatively, an every third day dosing regimen targeted to maintain FVIII trough levels $\geq 1\%$ may be employed. Adjust dose based on the patient's clinical response.
Perioperative management Congenital Hemophilia A	<u>Minor</u> Circulating Factor VIII required (% of normal) (60-100%) = 30-50 IU/ kg –Single dose within one hour of the operation. Repeat after 12- 24 hours for optional additional dosing as needed to control bleeding. <u>Major</u> Circulating Factor VIII required (% of normal) (80-120%) = Preoperative: 40-60 IU/ kg to achieve 100% activity. Followed by a repeat dose every 8-24 hours (every 6 to 24 hours for patients under age of 6) postoperatively until healing is complete.

Adynovate

Indication	Dose
On-demand treatment and control of bleeding episodes Congenital Hemophilia A	Dose (IU) = Body Weight (kg) x Desired factor VIII rise (IU/dL or % of normal) x 0.5 (IU/kg per IU/dL) <u>Minor</u> Target Factor VIII level (IU/dL or % of normal) (20-40%) = 10-20 IU/kg -Repeat every 12-24 hours until the bleeding episode is resolved <u>Moderate</u> Target Factor VIII level (IU/dL or % of normal) (30-60%) = 15-30 IU/kg - Repeat every 12-24 hours until the bleeding episode is resolved <u>Major</u> Target Factor VIII level (IU/dL or % of normal) (60-100%) = 30-50 IU/kg - Repeat every 8-24 hours until the bleeding episode is resolved.

FACTOR VIII_HEMOPHILIA PRODUCTS - Prior Auth Criteria

Proprietary Information. Restricted Access – Do not disseminate or copy without approval.

©2023, Magellan Rx Management

Indication	Dose
Perioperative management Congenital Hemophilia A	<u>Minor</u> Target Factor VIII required (% of normal) (60-100%) = 30-50 IU/ kg –Single dose within one hour of the operation. Repeat after 24 hours, if necessary, single dose or repeat as needed until bleeding is resolved. <u>Major</u> Target Factor VIII required (% of normal) (80-120%) (pre- and post- operative) = 40-60 IU/ kg within 1 hour of the operation to achieve 100% activity. Repeat dose every 8-24 hours (every 6 to 24 hours for patients under age of 12) to maintain FVIII activity within the target range and continue until adequate wound healing.
Routine prophylaxis Congenital Hemophilia A	Administer 40-50 IU per kg body weight 2 times per week in children and adults (12 years and older). Administer 55 IU per kg body weight 2 times per week in children (<12 years) with a maximum of 70 IU per kg. Adjust the dose based on the patient's clinical response.

Afstyla

Indication	Dose
On-demand treatment and control of bleeding episodes Congenital Hemophilia A	Dose (IU) = Body Weight (kg) x Desired factor VIII rise (IU/dL or % of normal) x 0.5 (IU/kg per IU/dL) <u>Minor</u> Target Factor VIII level (IU/dL or % of normal) 20-40% -Repeat every 12-24 hours until the bleeding episode is resolved. <u>Moderate</u> Target Factor VIII level (IU/dL or % of normal) 30-60%- Repeat every 12-24 hours until the bleeding episode is resolved. <u>Major</u> Target Factor VIII level (IU/dL or % of normal) 60-100%- Repeat every 8-24 hours until the bleeding episode is resolved.
Perioperative management Congenital Hemophilia A	<u>Minor</u> Target Factor VIII level (IU/dL or % of normal) 30-60%- Repeat every 24 hours, for at least one day, until healing is achieved. <u>Major</u> Target Factor VIII level (IU/dL or % of normal) 80-100%- Repeat every 8-24 hours until adequate wound healing, then continue for at least another 7 days to maintain a Factor VIII activity of 30-60% (IU/dL).
Routine prophylaxis Congenital Hemophilia A	Adults and adolescents (≥ 12 yrs old): Administer 20-50 IU per kg body weight 2 to 3 times per week. Adjust the dose based on the patient's clinical response. Children (<12 yrs old): Administer 30-50 IU per kg body weight 2 to 3 times per week. Adjust the dose based on the patient's clinical response.

Altuviiiio

Indication	Dose
On-demand treatment and control of bleeding episodes Congenital Hemophilia A	<u>Minor/Moderate</u> Single dose of 50 IU/kg. For minor and moderate bleeding episodes occurring within 2 to 3 days after a prophylactic dose, a lower dose of 30 IU/kg dose may be used. Additional doses of 30 or 50 IU/kg every 2 to 3 days may be considered. <u>Major</u> Single dose of 50 IU/kg. Additional doses of 30 or 50 IU/kg every 2 to 3 days can be considered. <i>Note: For resumption of prophylaxis (if applicable) after treatment of a bleed, it is recommended to allow an interval of at least 72 hours between the last 50 IU/kg dose for treatment of a bleed and resuming prophylaxis dosing. Thereafter, prophylaxis can be continued as usual on the patient's regular schedule.</i>
Perioperative management Congenital Hemophilia A	<u>Minor</u> Single dose of 50 IU/kg. An additional dose of 30 or 50 IU/kg after 2 to 3 days may be considered. <u>Major</u> Single dose of 50 IU/kg. Additional doses of 30 or 50 IU/kg every 2 to 3 days may be administered as clinically needed for perioperative management.
Routine prophylaxis Congenital Hemophilia A	The recommended dosing for routine prophylaxis for adults and children is 50 IU/kg of Altuviiiio administered once weekly.
– For the dose of 50 IU/kg, the expected in vivo peak increase in Factor VIII level expressed as IU/dL (or % of normal) is estimated using the following formula: – Estimated Increment of Factor VIII (IU/dL or % of normal) = 50 IU/kg x 2 (IU/dL per IU/kg) – To achieve a specific target Factor VIII activity level, use the following formula: Dosage (IU) = Body Weight (kg) x Desired Factor VIII Increase (IU/dL or % normal) x 0.5 (IU/kg per IU/dL).	

Eloctate

Indication	Dose
On-demand treatment and control of bleeding episodes Congenital Hemophilia A	Dose (IU/kg) = Desired factor VIII rise (IU/dL or % of normal) x 0.5 (IU/kg per IU/dL) <u>Minor and Moderate</u> Circulating Factor VIII required (% of normal) (40-60%) = 20-30 IU/ kg -Repeat every 24-48 hours as needed (every 12 to 24 hours for patients under age of 6). Continue until the bleeding episode is resolved. <u>Major</u> Circulating Factor VIII required (% of normal) (80-100%) = 40-50 IU/ kg - Repeat every 12-24 hours as needed (every 8 to 24 hours for patients under age of 6). Continue until the bleeding episode is resolved (approximately 7-10 days).

Indication	Dose
Routine prophylaxis Congenital Hemophilia A	Adults: The recommended starting regimen is 50 IU/kg administered every 4 days. The regimen may be adjusted based on patient response with dosing in the range of 25-65 IU/kg at 3–5-day intervals. Children < 6 years of age: The recommended starting regimen is 50 IU/kg administered twice weekly. The regimen may be adjusted based on patient response with dosing in the range of 25-65 IU/kg at 3–5-day intervals. More frequent or higher doses up to 80 IU/kg may be required.
Perioperative management Congenital Hemophilia A	<u>Minor</u> Circulating Factor VIII required (% of normal) (50-80%) = 25-40 IU/ kg -Repeat every 24 hours as needed (every 12 to 24 hours for patients under age of 6). Continue at least 1 day until healing is achieved. <u>Major</u> Circulating Factor VIII required (% of normal) (80-120%) = Preoperative: 40-60 IU/ kg – Followed by a repeat dose of 40-50 IU/kg after 8-24 hours (6 to 24 hours for patients under age of 6). Continue every 24 hours until adequate wound healing; then continue therapy for at least 7 days to maintain FVII activity within the target range.

Esperoct

Indication	Dose			
On-demand treatment and control of bleeding episodes Congenital Hemophilia A	One IU of Factor VIII activity corresponds to the quantity of Factor VIII in one milliliter of normal human plasma. The calculation of the required dosage of Factor VIII is based on the empirical finding that one IU of Factor VIII per kg body weight raises the plasma Factor VIII activity by two IU/dL.			
	To achieve a specific target Factor VIII activity level, use the following formula: Dosage (IU) = Body Weight (kg) × Desired Factor VIII Increase (IU/dL or % normal) × 0.5 ; OR			
	Type of bleeding	Adolescents/Adults ≥12 years Dose (IU/kg)	Children <12 years Dose (IU/kg)	Additional doses
	Minor Early hemarthrosis, mild muscle bleeding, or oral bleeding	40	65	One dose should be sufficient
	Moderate More extensive hemarthrosis, muscle bleeding, or hematoma	40	65	An additional dose may be administered after 24 hours
	Major Life- or limb-threatening hemorrhages, gastro- intestinal bleeding, intracranial, intra-abdominal or intrathoracic bleeding, fractures	50	65	Additional dose(s) may be administered approximately every 24 hours
Routine prophylaxis Congenital Hemophilia A	– Adults and adolescents (≥ 12 years): The recommended starting dose is 50 IU per kg body weight every 4 days. This regimen may be individually adjusted to less or more frequent dosing based on bleeding episodes. – Children (< 12 years): A dose of 65 IU per kg body weight twice weekly. This regimen may be individually adjusted to less or more frequent dosing based on bleeding episodes.			

FACTOR VIII_HEMOPHILIA PRODUCTS - Prior Auth Criteria

Proprietary Information. Restricted Access – Do not disseminate or copy without approval.

©2023, Magellan Rx Management

Indication	Dose			
Perioperative management Congenital Hemophilia A	<i>To achieve a specific target Factor VIII activity level, use the following formula: Dosage (IU) = Body Weight (kg) × Desired Factor VIII Increase (IU/dL or % normal) × 0.5 ; OR</i>			
	Type of surgery	Adolescents/Adults ≥12 years Dose (IU/kg)	Children <12 years Dose (IU/kg)	Additional doses
	Minor Including tooth extraction	50	65	Additional dose(s) can be given after 24 hours if necessary
	Major Intracranial, intra-abdominal, intrathoracic, or joint replacement surgery	50	65	Additional doses can be given every 24 hours for the first week and then approximately every 48 hours until wound healing has occurred

Hemofil M

Indication	Dose
On-demand treatment and control of bleeding episodes Congenital Hemophilia A	Dose (IU/kg) = Desired factor VIII rise (IU/dL or % of normal) x 0.5 (IU/kg per IU/dL) <u>Early hemarthrosis or muscle bleed or oral bleed</u> Circulating Factor VIII required (% of normal) (20-40%) = - Begin infusion every 12 to 24 hours for one-three days until the bleeding episode as indicated by pain is resolved or healing is achieved. <u>More extensive hemarthrosis, muscle bleed, or hematoma</u> Circulating Factor VIII required (% of normal) (30-60%) = Repeat every 12-24 hours for usually three days or more until pain and disability are resolved. <u>Life threatening bleeds such as head injury, throat bleed, severe abdominal pain</u> Circulating Factor VIII Required (% of normal) (60-100%) = Repeat every 8-24 hours until the bleeding threat is resolved.
Perioperative management Congenital Hemophilia A	<u>Minor</u> Circulating Factor VIII required (% of normal) (60-80%) A single infusion plus oral antifibrinolytic therapy within one hour is sufficient in approximately 70% of cases. <u>Major</u> Circulating Factor VIII required (% of normal) (80-100% pre- and post-operative): Repeat dose every 8-24 hours depending on state of healing.

Jivi

Indication	Dose
On-demand treatment and control of bleeding episodes Congenital Hemophilia A	Dose (IU/kg) = Desired factor VIII rise (IU/dL or % of normal) x reciprocal of expected recovery (or observed recovery, if available) (e.g., 0.5 for a recovery of 2 IU/dL per IU/kg) <u>Minor</u> Circulating Factor VIII required (% of normal) (20-40%) – 10-20IU/kg repeat dose every 24-48 hours until bleed resolves <u>Moderate</u>

Indication	Dose
	Circulating Factor VIII required (% of normal) (30-60%) – 15-30IU/kg repeat dose every 24-48 hours until bleed resolves <u>Major</u> Circulating Factor VIII Required (% of normal) (60-100%) – 30-50IU/kg repeat dose every 8-24 hours until bleed resolves
Perioperative management Congenital Hemophilia A	<u>Minor</u> Circulating Factor VIII required (% of normal) (30-60%) – 15-30IU/kg repeat dose every 24 hours for at least 1 day until healing is achieved <u>Major</u> Circulating Factor VIII required (% of normal) (80-100%) – 40-50IU/kg repeat dose every 12-24 hours until adequate wound healing is complete, then continue therapy for at least another 7 days to maintain Factor VIII activity of 30–60% (IU/dL)
Routine prophylaxis Congenital Hemophilia A	The recommended initial regimen is 30–40 IU/kg twice weekly. Based on the bleeding episodes, the regimen may be adjusted to 45–60 IU/kg every 5 days or may be further individually adjusted to less or more frequent dosing.

Koate/Koate DVI

Indication	Dose
On-demand treatment and control of bleeding episodes Congenital Hemophilia A	Dose (IU/kg) = Desired factor VIII rise (IU/dL or % of normal) x 0.5 (IU/kg per IU/dL) <u>Mild</u> Circulating Factor VIII required (% of normal) (20%) = 10 IU/kg- Therapy need not be repeated unless there is evidence of further bleeding. <u>Moderate</u> Circulating Factor VIII required (% of normal) (30-50%) = 15-25 IU/kg - If further therapy is required, repeated doses of 10-15 IU per kg every 8-12 hours may be given. <u>Severe</u> Circulating Factor VIII Required (% of normal) (80-100%) =40-50 IU/kg – followed by a maintenance dose of 20-25 IU per kg every 8-12 hours.
Routine prophylaxis Hemophilia A §	25-40 IU/kg three times weekly or 15-30 IU/kg three times weekly. Adjust dosing regimen based on individual response.
Perioperative management Congenital Hemophilia A	For major surgical procedures, the Factor VIII level should be raised to approximately 100% by giving a preoperative dose of 50 IU/kg. The Factor VIII level should be checked to assure that the expected level is achieved before the patient goes to surgery. To maintain hemostatic levels, repeat infusions may be necessary every 6 to 12 hours initially, and for a total of 10 to 14 days until healing is complete. The intensity of Factor VIII replacement therapy required depends on the type of surgery and postoperative regimen employed. For minor

Indication	Dose
	surgical procedures, less intensive treatment schedules may provide adequate hemostasis.

Kogenate FS

Indication	Dose
On-demand treatment and control of bleeding episodes Congenital Hemophilia A	Dose (IU/kg) = Desired factor VIII rise (IU/dL or % of normal) x 0.5 (IU/kg per IU/dL) <u>Minor</u> Circulating Factor VIII required (% of normal) (20-40%) = 10-20 IU/ kg -Repeat dose if there is evidence of further bleeding and continue until the bleeding episode is resolved. <u>Moderate</u> Circulating Factor VIII required (% of normal) (30-60%) = 15-30 IU/ kg - Repeat every 12-24 hours as needed. Continue until the bleeding episode is resolved. <u>Major</u> Circulating Factor VIII Required (% of normal) (80-100%) = Initial: 40-50 IU/ kg; Repeat 20-25 IU/kg every 8-12 hours until the bleeding episode is resolved.
Routine prophylaxis Congenital Hemophilia A	<u>Routine Prophylaxis in Adults</u> 25 units per kg of body weight three times per week. <u>Routine Prophylaxis in Children</u> 25 IU/kg of body weight every other day.
Perioperative management Congenital Hemophilia A	<u>Minor</u> Circulating Factor VIII required (% of normal) (30-60%) = 15-30 IU/ kg – Repeat every 12- 24 hours until bleeding is resolved. <u>Major</u> Circulating Factor VIII required (% of normal) (100%) = Preoperative: 50 IU/ kg to achieve 100% activity. Followed by a repeat dose every 6-12 hours to keep FVIII activity in desired range. Continue until healing is complete.

Kovaltry

Indication	Dose
On-demand treatment and control of bleeding episodes Congenital Hemophilia A	- Required dose (IU) = body weight (kg) x desired Factor VIII rise (% of normal or IU/dL) x reciprocal of expected/observed recovery (e.g., 0.5 for a recovery of 2 IU/dL per IU/kg) - Estimated Increment of Factor VIII (IU/dL or % of normal) = [Total Dose (IU)/body weight (kg)] x 2 (IU/dL per IU/kg) <u>Minor</u> (Early hemarthrosis, minor muscle, oral bleeds) Factor VIII level required (IU/dL or % of normal): 20-40 – repeat every 12-24 hours at least 1 day, until bleeding episode as indicated by pain is resolved or healing is achieved.

FACTOR VIII_HEMOPHILIA PRODUCTS - Prior Auth Criteria

Proprietary Information. Restricted Access – Do not disseminate or copy without approval.

©2023, Magellan Rx Management

Indication	Dose
	<u>Moderate</u> (More extensive hemarthrosis, muscle bleeding, or hematoma) Factor VIII level required (IU/dL or % of normal): 30-60 – repeat every 12-24 hours for 3 to 4 days or more until pain and acute disability are resolved. <u>Major</u> (Intracranial, intra-abdominal or intrathoracic hemorrhages, gastrointestinal bleeding, central nervous system bleeding, bleeding in the retropharyngeal or retroperitoneal spaces, or iliopsoas sheath, life or limb threatening hemorrhage) Factor VIII level required (IU/dL or % of normal): 60-100 – repeat every 8-24 hours until bleeding is resolved.
Routine prophylaxis Congenital Hemophilia A	Individualize the patient's dose based on clinical response: - Adults and adolescents: 20 to 40 IU of KOVALTRY per kg of body weight two or three times per week. - Children ≤12 years old: 25 to 50 IU of KOVALTRY per kg body weight twice weekly, three times weekly, or every other day according to individual requirements
Perioperative management Congenital Hemophilia A	<u>Minor</u> (Such as tooth extraction) Factor VIII level required (IU/dL or % of normal): 30-60 (pre- and post-operative) – repeat every 24 hours at least 1 day until healing is achieved. <u>Major</u> (Such as intracranial, intraabdominal, intrathoracic, or joint replacement surgery) Factor VIII level required (IU/dL or % of normal): 80-100 (pre- and post-operative) – repeat every 8-24 hours until adequate wound healing is complete, then continue therapy for at least another 7 days to maintain Factor VIII activity of 30-60% (IU/dL).

Novoeight

Indication	Dose
On-demand treatment and control of bleeding episodes Congenital Hemophilia A	Dose (IU/kg) = Desired factor VIII rise (IU/dL or % of normal) x 0.5 (IU/kg per IU/dL) <u>Minor</u> Circulating Factor VIII required (% of normal) (20-40%), every 12 – 24 hours for at least 1 day until the bleeding episode is resolved <u>Moderate</u> Circulating Factor VIII required (% of normal) (30-60%), every 12 – 24 hours until pain and acute disability are resolved, approximately 3-4 days <u>Major</u> Circulating Factor VIII Required (% of normal) (60-100%), every 8 – 24 hours until resolution of bleed, approximately 7-10 days.

Indication	Dose
Perioperative management Hemophilia A	Dose (IU/kg) = Desired factor VIII rise (IU/dL or % of normal) x 0.5 (IU/kg per IU/dL) <u>Minor</u> Circulating Factor VIII required (% of normal) (30-60%) every 24 hours for at least 1 day until healing is achieved. <u>Major</u> Circulating Factor VIII required (% of normal) (80-100%) every 8 – 24 hours until adequate wound healing, then continue therapy for at least 7 days to maintain a factor VIII activity of 30 – 60% (IU/dL)
Routine prophylaxis Hemophilia A	<u>Adults and adolescents (>12 yrs):</u> 20-50 IU/kg three times weekly OR 20-40 IU/kg every other day <u>Children (<12 yrs):</u> 25-60 IU/kg three times weekly OR 25-50 IU/kg every other day

NUWIQ

Indication	Dose
On-demand treatment and control of bleeding episodes Congenital Hemophilia A	<u>Dose</u> Required IU = body weight (kg) x desired Factor VIII rise (%) (IU/dL) x 0.5 (IU/kg per IU/dL) Expected Factor VIII rise (% of normal) = 2 x administered IU/body weight (kg) <u>Minor</u> Required peak post-infusion Factor VIII activity (% of normal or IU/dL): 20-40 every 12 – 24 hours for at least 1 day, until the bleeding episode is resolved <u>Moderate to Major</u> Required peak post-infusion Factor VIII activity (% of normal or IU/dL): 30-60 every 12 – 24 hours for 3-4 days or more until the bleeding episode is resolved <u>Life-threatening</u> Required peak post-infusion Factor VIII activity (% of normal or IU/dL): 60-100 every 8 – 24 hours bleeding risk is resolved
Routine prophylaxis Congenital Hemophilia A	<u>Dose</u> Required IU = body weight (kg) x desired Factor VIII rise (%) (IU/dL) x 0.5 (IU/kg per IU/dL) Expected Factor VIII rise (% of normal) = 2 x administered IU/body weight (kg) <u>Adolescents (12-17 years) and adults</u> 30 – 40 IU/kg every other day <u>Children (2-11 years)</u> 30 – 50 IU/kg every other day or three times per week

Indication	Dose
Perioperative management Congenital Hemophilia A	<u>Dose</u> Required IU = body weight (kg) x desired Factor VIII rise (%) (IU/dL) x 0.5 (IU/kg per IU/dL) Expected Factor VIII rise (% of normal) = 2 x administered IU/body weight (kg) <u>Minor</u> Required peak post-infusion Factor VIII activity (% of normal or IU/dL): 30-60 (pre- and post-operative) every 24 hours for at least 1 day until healing is achieved <u>Major</u> Required peak post-infusion Factor VIII activity (% of normal or IU/dL): 80-100 (pre- and post-operative) every 8 - 24 hours until adequate wound healing, then continue therapy for at least another 7 days to maintain Factor VIII activity of 30% to 60% (IU/dL)

Obizur

Indication	Dose
On-demand treatment and control of bleeding episodes Acquired Hemophilia A	<u>Minor and Moderate</u> Loading dose: 200IU/kg; Maintenance dose: Titrate to maintain recommended FVIII trough levels at 50-100 IU/dL every 4 to 12 hours <u>Major</u> Loading dose: 200 IU/kg; Maintenance dose: Titrate to maintain recommended FVIII trough levels at 100-200 (to treat an acute bleed), then 50-100 IU/dL (after acute bleed is controlled) every 4 to 12 hours

Recombinate

Indication	Dose
On-demand treatment and control of bleeding episodes Congenital Hemophilia A	Dose (IU/kg) = Desired factor VIII rise (IU/dL or % of normal) x 0.5 (IU/kg per IU/dL) <u>Early hemarthrosis or muscle bleed or oral bleed</u> Circulating Factor VIII required (% of normal) (20-40%) - Begin infusion every 12 to 24 hours for one-three days until the bleeding episode as indicated by pain is resolved or healing is achieved. <u>More extensive hemarthrosis, muscle bleed, or hematoma</u> Circulating Factor VIII required (% of normal) (30-60%) - Repeat every 12-24 hours for usually three days or more until pain and disability are resolved. <u>Life threatening bleeds such as head injury, throat bleed, severe abdominal pain</u> Circulating Factor VIII Required (% of normal) (60-100%) - Repeat every 8-24 hours until the bleeding threat is resolved.
Routine prophylaxis Hemophilia A §	25-40 IU/kg three times weekly or 15-30 IU/kg three times weekly. Adjust dosing regimen based on individual response.

Indication	Dose
Perioperative management Congenital Hemophilia A	<u>Minor</u> Circulating Factor VIII required (% of normal) (60-80%) - A single infusion plus oral antifibrinolytic therapy within one hour is sufficient in approximately 70% of cases. <u>Major</u> Circulating Factor VIII required (% of normal) (80-100% pre- and post-operative) - Repeat dose every 8-24 hours depending on state of healing.

Xyntha/Xyntha Solofuse

Indication	Dose
On-demand treatment and control of bleeding episodes Congenital Hemophilia A	Dose (IU/kg) = Desired factor VIII rise (IU/dL or % of normal) x 0.5 (IU/kg per IU/dL) <u>Minor</u> Circulating Factor VIII required (% of normal) (20-40%) - Repeat dose every 12-24 hours for least 1 day, depending upon the severity of the bleeding episode. <u>Moderate</u> Circulating Factor VIII required (% of normal) (30-60%) - Repeat every 12-24 hours as needed. Continue for 3-4 days or until adequate local hemostasis is achieved. <u>Major</u> Circulating Factor VIII Required (% of normal) (60-100%) - Repeat every 8-24 hours until bleeding is resolved.
Perioperative management Congenital Hemophilia A	<u>Minor</u> Circulating Factor VIII required (% of normal) (30-60%) - Repeat every 12-24 hours. Continue for 3-4 days or until adequate local hemostasis is achieved. For tooth extraction, a single infusion plus oral antifibrinolytic therapy within 1 hour may be sufficient. <u>Major</u> Circulating Factor VIII required (% of normal) (60-100%) - Repeat every 8-24 hours. Continue until threat is resolved, or in the case of surgery, until adequate local hemostasis and wound healing are achieved.
Routine prophylaxis Hemophilia A	<u>Adults and adolescents (>12 years):</u> The recommended starting regimen is 30 IU/kg of Xyntha administered 3 times weekly. <u>Children (<12 years):</u> The recommended starting regimen is 25 IU/kg of Xyntha administered every other day. More frequent or higher doses may be required in children <12 years of age to account for the higher clearance in this age group. Note: Adjust the dosing regimen (dose or frequency) based on the patient's clinical response.

§ Utrecht and/or Malmö protocols used as basis for dosing

VII. Billing Code/Availability Information

HCPSC Code & NDC:

Drug	Manufacturer	HCPSC Codes	1 Billable Unit Equiv.	Vial Size	NDC
Advate	Baxalta US Inc	J7192	1 IU	250 units	00944-3051-02
				500 units	00944-3052-02
				1000 units	00944-3053-02
				1500 units	00944-3054-02
				2000 units	00944-3045-10
				3000 units	00944-3046-10
				4000 units	00944-3047-10
Kogenate FS	Bayer HealthCare LLC	J7192	1 IU	250 units	00026-3782-25
				500 units	00026-3783-35
				1000 units	00026-3785-55
				2000 units	00026-3786-65
				3000 units	00026-3787-75
Recombinate	Baxalta US Inc.	J7192	1 IU	220-400 units	00944-2841-10
				401-800 units	00944-2842-10
				801-1240 units	00944-2843-10
				1241-1800 units	00944-2844-10
				1801-2400 units	00944-2845-10
Kovaltry	Bayer HealthCare LLC	J7211	1 IU	250 units	00026-3821-25
				500 units	00026-3822-25
				1000 units	00026-3824-25
				2000 units	00026-3826-50
				3000 units	00026-3828-50
Eloctate	Bioverativ Therapeutics Inc.	J7205	1 IU	250 units	71104-0801-01
				500 units	71104-0802-01
				750 units	71104-0803-01
				1000 units	71104-0804-01
				1500 units	71104-0805-01
				2000 units	71104-0806-01
				3000 units	71104-0807-01
				4000 units	71104-0808-01
				5000 units	71104-0809-01
				6000 units	71104-0810-01
Koate/Koate DVI	Grifols Therapeutics Inc	J7190	1 IU	250 units	76125-0250-20
					76125-0253-25
					76125-0256-20
					76125-0257-25
					76125-0258-02
					76125-0259-02
				500 units	76125-0661-02
					76125-0662-50
					76125-0663-50
				1000 units	76125-0665-02
					76125-0667-30
					76125-0668-30
					76125-0672-50
					76125-0674-10

FACTOR VIII_HEMOPHILIA PRODUCTS - Prior Auth Criteria

Proprietary Information. Restricted Access – Do not disseminate or copy without approval.

©2023, Magellan Rx Management

					76125-0675-12 76125-0676-50 76125-0678-10 76125-0679-12
Hemofil M	Takeda Pharmaceuticals USA, Inc.	J7190	1 IU	250 units	00944-3940-02
				500 units	00944-3942-02
				1000 units	00944-3944-02
				1700 units	00944-3946-02
Novoeight	Novo Nordisk Inc.	J7182	1 IU	250 units	00169-7825-01
				500 units	00169-7850-01
				1000 units	00169-7810-01
				1500 units	00169-7815-01
				2000 units	00169-7820-01
				3000 units	00169-7830-01
Nuwiq	Octapharma AB	J7209	1 IU	250 units	68982-0140-01
				500 units	68982-0142-01
				1000 units	68982-0144-01
				1500 units	68982-0154-01
				2000 units	68982-0146-01
				2500 units	68982-0148-01
				3000 units	68982-0150-01
				4000 units	68982-0152-01
Obizur	Baxalta US Inc.	J7188	1 IU	500 units	00944-5001-xx
Xyntha/Xyntha Solofuse	Wyeth Pharmaceuticals LLC	J7185	1 IU	250 units	58394-0012-01 58394-0022-03
				500 units	58394-0013-01 58394-0023-03
				1000 units	58394-0014-01 58394-0024-03
				2000 units	58394-0015-01 58394-0025-03
				3000 units	58394-0016-03
AfstylA	CSL Behring, LLC	J7210	1 IU	250 units	69911-0474-02
				500 units	69911-0475-02
				1000 units	69911-0476-02
				1500 units	69911-0480-02
				2000 units	69911-0477-02
				2500 units	69911-0481-02
Adynovate	Baxalta US Inc.	J7207	1 IU	3000 units	69911-0478-02
				250 units	00944-4622-01
				500 units	00944-4623-01
				750 units	00944-4626-01
				1000 units	00944-4624-01
				1500 units	00944-4627-01
				2000 units	00944-4625-01
Jivi	Bayer HealthCare LLC	J7208	1 IU	3000 units	00944-4628-01
				500 units	00026-3942-25
				1000 units	00026-3944-25
				2000 units	00026-3946-25
Esperoct	Novo Nordisk Inc.	J7204	1 IU	3000 units	00026-3948-25
				500 units	00169-8500-01
				1000 units	00169-8100-01
				1500 units	00169-8150-01

FACTOR VIII_HEMOPHILIA PRODUCTS - Prior Auth Criteria

Proprietary Information. Restricted Access – Do not disseminate or copy without approval.

©2023, Magellan Rx Management

				2000 units	00169-8200-01
				3000 units	00169-8300-01
Altuviiiio	Bioverativ Therapeutics Inc.	J7199	N/A	250 units	71104-0978-01
				500 units	71104-0979-01
				750 units	71104-0980-01
				1000 units	71104-0981-01
				2000 units	71104-0982-01
				3000 units	71104-0983-01
				4000 units	71104-0984-01

VIII. References

1. Advate [package insert]. Lexington, MA; Baxalta US Inc. March 2023. Accessed May 2023.
2. Afstylia [package insert]. Kankakee, IL; CSL Behring, LLC; April 2021. Accessed May 2023.
3. Eloctate [package insert]. Waltham, MA; Bioverativ Therapeutics Inc.; December 2020. Accessed May 2023.
4. Hemofil M [package insert]. Lexington, MA; Takeda Pharmaceuticals USA, Inc. March 2023. Accessed May 2023.
5. Koate/Koate DVI [package insert]. Research Triangle Park, NC; Grifols Therapeutics Inc.; January 2022. Accessed May 2023.
6. Kogenate FS [package insert]. Whippany, NJ. Bayer HealthCare LLC; December 2019. Accessed May 2023.
7. Novoeight [package insert]. Bagsvaerd, Denmark; Novo Nordisk Inc.; July 2020. Accessed May 2023.
8. NUWIQ [package insert]. Elersvagen, Sweden; Octapharma AB; June 2021. Accessed May 2023.
9. Obizur [package insert]. Lexington, MA; Baxalta US Inc. March 2023. Accessed May 2023.
10. Recombinate [package insert]. Lexington, MA; Baxalta US Inc. March 2023. Accessed April 2022.
11. Xyntha & Xyntha Solofuse [package insert]. Philadelphia, PA; Wyeth Pharmaceuticals LLC; July 2022. Accessed May 2023.
12. Adynovate [package insert]. Lexington, MA; Baxalta US Inc.; March 2023. Accessed May 2023.
13. Kovaltry [package insert]. Whippany, NJ; Bayer HealthCare LLC; December 2022. Accessed May 2023.
14. Jivi [package insert]. Whippany, NJ; Bayer HealthCare LLC; August 2018. Accessed May 2023.
15. Altuviiiio [package insert]. Waltham, MA; Bioverativ Therapeutics Inc.; March 2023. Accessed May 2023.
16. MASAC RECOMMENDATIONS CONCERNING PRODUCTS LICENSED FOR THE TREATMENT OF HEMOPHILIA AND OTHER BLEEDING DISORDERS. Revised August 2020 National Hemophilia Foundation. MASAC Document #263; August 2020. Available at: <http://www.hemophilia.org>. Accessed May 2023.
17. Guidelines for the Management of Hemophilia. 3rd Edition. World Federation of Hemophilia. 2020. Available at: <https://www1.wfh.org/publications/files/pdf-1863.pdf>. Accessed May 2023.

18. Annual Review of Factor Replacement Products. Oklahoma Health Care Authority Review Board. Updated April 2016. Accessed May 2023.
19. Graham A1, Jaworski K. Pharmacokinetic analysis of anti-hemophilic factor in the obese patient. *Haemophilia*. 2014 Mar;20(2):226-9.
20. Croteau SE1, Neufeld EJ. Transition considerations for extended half-life factor products. *Haemophilia*. 2015 May;21(3):285-8.
21. Mingot-Castellano, et al. Application of Pharmacokinetics Programs in Optimization of Haemostatic Treatment in Severe Hemophilia a Patients: Changes in Consumption, Clinical Outcomes and Quality of Life. *Blood*. 2014 December; 124 (21).
22. MASAC RECOMMENDATION CONCERNING PROPHYLAXIS. 2016 National Hemophilia Foundation. MASAC Document #241; February 2016. Available at: <http://www.hemophilia.org>. Accessed May 2023.
23. Esperoct [package insert]. Plainsboro, NJ; Novo Nordisk Inc.; August 2022. Accessed May 2023.
24. First Coast Service Options, Inc. Local Coverage Article: Billing and Coding: Hemophilia Clotting Factors (A56482). Centers for Medicare & Medicaid Services Inc. Updated on 10/28/2022 with effective date 10/01/2022. Accessed May 2023.
25. Palmetto GBA. Local Coverage Article: Billing and Coding: Guidance for Anti-Inhibitor Coagulant Complex (AICC) National Coverage Determination (NCD) 110.3 (A56065). Centers for Medicare & Medicaid Services Inc. Updated on 11/14/2022 with effective date 11/24/2022. Accessed May 2023.
26. Novitas Solutions, Inc. Local Coverage Article: Billing and Coding: Hemophilia Factor Products (A56433). Centers for Medicare & Medicaid Services Inc. Updated on 10/14/2022 with effective date 10/01/2022. Accessed May 2023.

Appendix 1 – Covered Diagnosis Codes

Obizur

ICD-10	ICD-10 Description
D68.311	Acquired hemophilia

Advate, Eloctate, Hemofil M, Koate-DVI, Kogenate FS, Recombinate, Xyntha/ Xyntha Solofuse, Novoeight. NUWIQ, Adynovate, Kovaltry, Afstylia, Jivi, and Altuviiiio

ICD-10	ICD-10 Description
D66	Hereditary factor VIII deficiency

Appendix 2 – Centers for Medicare and Medicaid Services (CMS)

Medicare coverage for outpatient (Part B) drugs is outlined in the Medicare Benefit Policy Manual (Pub. 100-2), Chapter 15, §50 Drugs and Biologicals. In addition, National Coverage Determination (NCD), Local Coverage Determinations (LCDs), and Local Coverage Articles (LCAs) may exist and compliance with these policies is required where applicable. They can be found at:

<https://www.cms.gov/medicare-coverage-database/search.aspx>. Additional indications may be covered at the discretion of the health plan.

Medicare Part B Covered Diagnosis Codes (applicable to existing NCD/LCD/LCA):

Jurisdiction(s): N	NCD/LCD Document (s): A56482
https://www.cms.gov/medicare-coverage-database/new-search/search-results.aspx?keyword=a56482&areaId=all&docType=NCA%2CCAL%2CNCD%2CMEDCAC%2CTA%2CMCD%2C6%2C3%2C5%2C1%2CF%2CP	

Jurisdiction(s): J,M	NCD/LCD Document (s): A56065
https://www.cms.gov/medicare-coverage-database/new-search/search-results.aspx?keyword=a56065&areaId=all&docType=NCA%2CCAL%2CNCD%2CMEDCAC%2CTA%2CMCD%2C6%2C3%2C5%2C1%2CF%2CP	

Jurisdiction(s): H,L	NCD/LCD Document (s): A56433
https://www.cms.gov/medicare-coverage-database/new-search/search-results.aspx?keyword=a56433&areaId=all&docType=NCA%2CCAL%2CNCD%2CMEDCAC%2CTA%2CMCD%2C6%2C3%2C5%2C1%2CF%2CP	

Medicare Part B Administrative Contractor (MAC) Jurisdictions		
Jurisdiction	Applicable State/US Territory	Contractor
E (1)	CA, HI, NV, AS, GU, CNMI	Noridian Healthcare Solutions, LLC
F (2 & 3)	AK, WA, OR, ID, ND, SD, MT, WY, UT, AZ	Noridian Healthcare Solutions, LLC
5	KS, NE, IA, MO	Wisconsin Physicians Service Insurance Corp (WPS)
6	MN, WI, IL	National Government Services, Inc. (NGS)
H (4 & 7)	LA, AR, MS, TX, OK, CO, NM	Novitas Solutions, Inc.
8	MI, IN	Wisconsin Physicians Service Insurance Corp (WPS)
N (9)	FL, PR, VI	First Coast Service Options, Inc.
J (10)	TN, GA, AL	Palmetto GBA, LLC
M (11)	NC, SC, WV, VA (excluding below)	Palmetto GBA, LLC
L (12)	DE, MD, PA, NJ, DC (includes Arlington & Fairfax counties and the city of Alexandria in VA)	Novitas Solutions, Inc.
K (13 & 14)	NY, CT, MA, RI, VT, ME, NH	National Government Services, Inc. (NGS)
15	KY, OH	CGS Administrators, LLC

PreferredOne Community Health Plan Nondiscrimination Notice

PreferredOne Community Health Plan (“PCHP”) complies with applicable Federal civil rights laws and does not discriminate on the basis of race, color, national origin, age, disability, or sex. PCHP does not exclude people or treat them differently because of race, color, national origin, age, disability, or sex.

PCHP:

Provides free aids and services to people with disabilities to communicate effectively with us, such as:

- Qualified sign language interpreters
- Written information in other formats (large print, audio, accessible electronic formats, other formats)

Provides free language services to people whose primary language is not English, such as:

- Qualified interpreters
- Information written in other languages

If you need these services, contact a Grievance Specialist.

If you believe that PCHP has failed to provide these services or discriminated in another way on the basis of race, color, national origin, age, disability, or sex, you can file a grievance with:

Grievance Specialist
PreferredOne Community Health Plan
PO Box 59052
Minneapolis, MN 55459-0052
Phone: 1.800.940.5049 (TTY: 763.847.4013)
Fax: 763.847.4010
customerservice@preferredone.com

You can file a grievance in person or by mail, fax, or email. If you need help filing a grievance, a Grievance Specialist is available to help you.

You can also file a civil rights complaint with the U.S. Department of Health and Human Services, Office for Civil Rights, electronically through the Office for Civil Rights Complaint Portal, available at <https://ocrportal.hhs.gov/ocr/portal/lobby.jsf>, or by mail or phone at:

U.S. Department of Health and Human Services
200 Independence Avenue, SW
Room 509F, HHH Building
Washington, D.C. 20201
1-800-368-1019, 800-537-7697 (TDD)

Complaint forms are available at <http://www.hhs.gov/ocr/office/file/index.html>.

Language Assistance Services

ATTENTION: If you do not speak English, language assistance services, free of charge, are available to you. Call 1.800.940.5049 (TTY: 763.847.4013).

ATENCIÓN: si habla español, tiene a su disposición servicios gratuitos de asistencia lingüística. Llame al 1.800.940.5049 (TTY: 763.847.4013).

LUS CEEV: Yog tias koj hais lus Hmoob, cov kev pab txog lus, muaj kev pab dawb rau koj. Hu rau 1.800.940.5049 (TTY: 763.847.4013).

XIYYEEFFANNAA: Afaan dubbattu Oroomiffa, taiaajiila qarqaarsa afaanii, kanfaltiidhaan ala, ni argama. Bilbilaa 1.800.940.5049 (TTY: 763.847.4013).

CHÚ Ý: Nếu bạn nói Tiếng Việt, có các dịch vụ hỗ trợ ngôn ngữ miễn phí dành cho bạn. Gọi số 1.800.940.5049 (TTY: 763.847.4013).

注意：如果您使用繁體中文，您可以免費獲得語言援助服務。請致電 1.800.940.5049 (TTY: 763.847.4013)。

ВНИМАНИЕ: Если вы говорите на русском языке, то вам доступны бесплатные услуги перевода. Звоните 1.800.940.5049 (телетайп: 763.847.4013).

ໂບດຊາບ: ຖ້າວ່າທ່ານເວົ້າພາສາລາວ, ການບໍລິການຊ່ວຍເຫຼືອດ້ານພາສາ, ໂດຍບໍ່ເສຍຄ່າ, ແມ່ນມີພ້ອມໃຫ້ທ່ານ. ໂທ 1.800.940.5049 (TTY: 763.847.4013).

ማስታወሻ፡ የሚናገሩት ቋንቋ አማርኛ ከሆነ የትርጉም እርዳታ ድርጅቶች፡ በነጻ ሊያግዝዎት ተዘጋጅተዋል፡ ወደ ሚከተለው ቁጥር ይደውሉ 1.800.940.5049 (መስማት ለተሳናቸው፡ 763.847.4013) .

ဟ်သ့ဟ်သး- နမၤကတိၤ ကသီၤ ကျိၣ်အယိၤ, နမၤန့ၣ် ကျိၣ်အတၢ်မၤစၢၤလၢ တလၢၣ်ဘျဉ်လၢၣ်စၢၤ နီၣ်တမံၤဘၣ်သ့န့ၣ်လီၤ. ကိး 1.800.940.5049 (TTY: 763.847.4013).

ACHTUNG: Wenn Sie Deutsch sprechen, stehen Ihnen kostenlos sprachliche Hilfsdienstleistungen zur Verfügung. Rufnummer: 1.800.940.5049 (TTY: 763.847.4013).

ប្រយ័ត្ន៖ បើសិនជាអ្នកនិយាយ ភាសាខ្មែរ, សេវាជំនួយផ្នែកភាសា ដោយមិនគិតល្បួល គឺអាចមានសំរាប់បំរើអ្នក។ ចរ ទូរស័ព្ទ 1.800.940.5049 (TTY: 763.847.4013)។

ملحوظة: إذا كنت تتحدث اذكر اللغة، فإن خدمات المساعدة اللغوية تتوافر لك بالمجان. اتصل برقم 1.800.940.5049 (رقم هاتف الصم والبكم: 763.847.4013).

ATTENTION : Si vous parlez français, des services d'aide linguistique vous sont proposés gratuitement. Appelez le 1.800.940.5049 (TTY: 763.847.4013).

주의: 한국어를 사용하시는 경우, 언어 지원 서비스를 무료로 이용하실 수 있습니다. 1.800.940.5049 (TTY: 763.847.4013), 번으로 전화해 주십시오.

PAUNAWA: Kung nagsasalita ka ng Tagalog, maaari kang gumamit ng mga serbisyo ng tulong sa wika nang walang bayad. Tumawag sa 1.800.940.5049 (TTY: 763.847.4013).

PreferredOne Insurance Company Nondiscrimination Notice

PreferredOne Insurance Company ("PIC") complies with applicable Federal civil rights laws and does not discriminate on the basis of race, color, national origin, age, disability, or sex. PIC does not exclude people or treat them differently because of race, color, national origin, age, disability, or sex.

PIC:

Provides free aids and services to people with disabilities to communicate effectively with us, such as:

- Qualified sign language interpreters
- Written information in other formats (large print, audio, accessible electronic formats, other formats)

Provides free language services to people whose primary language is not English, such as:

- Qualified interpreters
- Information written in other languages

If you need these services, contact a Grievance Specialist.

If you believe that PIC has failed to provide these services or discriminated in another way on the basis of race, color, national origin, age, disability, or sex, you can file a grievance with:

Grievance Specialist
PreferredOne Insurance Company
PO Box 59212
Minneapolis, MN 55459-0212
Phone: 1.800.940.5049 (TTY: 763.847.4013)
Fax: 763.847.4010
customerservice@preferredone.com

You can file a grievance in person or by mail, fax, or email. If you need help filing a grievance, a Grievance Specialist is available to help you.

You can also file a civil rights complaint with the U.S. Department of Health and Human Services, Office for Civil Rights, electronically through the Office for Civil Rights Complaint Portal, available at <https://ocrportal.hhs.gov/ocr/portal/lobby.jsf>, or by mail or phone at:

U.S. Department of Health and Human Services
200 Independence Avenue, SW
Room 509F, HHH Building
Washington, D.C. 20201
1-800-368-1019, 800-537-7697 (TDD)

Complaint forms are available at <http://www.hhs.gov/ocr/office/file/index.html>.

Language Assistance Services

ATTENTION: If you do not speak English, language assistance services, free of charge, are available to you. Call 1.800.940.5049 (TTY: 763.847.4013).

ATENCIÓN: si habla español, tiene a su disposición servicios gratuitos de asistencia lingüística. Llame al 1.800.940.5049 (TTY: 763.847.4013).

LUS CEEV: Yog tias koj hais lus Hmoob, cov kev pab txog lus, muaj kev pab dawb rau koj. Hu rau 1.800.940.5049 (TTY: 763.847.4013).

XIYEEFFANNAA: Afaan dubbattu Oroomiffa, tajaajila gargaarsa afaanii, kanfaltiidhaan ala, ni argama. Bilbilaa 1.800.940.5049 (TTY: 763.847.4013).

CHÚ Ý: Nếu bạn nói Tiếng Việt, có các dịch vụ hỗ trợ ngôn ngữ miễn phí dành cho bạn. Gọi số 1.800.940.5049 (TTY: 763.847.4013).

注意：如果您使用繁體中文，您可以免費獲得語言援助服務。請致電 1.800.940.5049 (TTY: 763.847.4013)。

ВНИМАНИЕ: Если вы говорите на русском языке, то вам доступны бесплатные услуги перевода. Звоните 1.800.940.5049 (телетайп: 763.847.4013).

បំពេញ: ប្រសិនបើ អ្នកនិយាយភាសាខ្មែរ, សេវាជំនួយភាសា ដោយមិនគិតថ្លៃ គឺអាចមានសំរាប់អ្នក។ ហៅ 1.800.940.5049 (TTY: 763.847.4013).

ማስታወሻ: የሚናገሩት ቋንቋ አማርኛ ከሆነ የትርጉም እርዳታ ድርጅቶች፡ በነጻ ሊያገኙበት ተዘጋጅተዋል፡ ወደ ሚከተለው ቁጥር ይደውሉ 1.800.940.5049 (ማስማት ለተሳናቸው: 763.847.4013) .

ဟံသာဝတီ: နမူနာတို့ ကညီ ကျိအသိ, နမူနာ ကျိအတိအကျတို့ တလက်တလက်စွာ နှိတ်တံ့သုန့်လိ။ ကိ: 1.800.940.5049 (TTY: 763.847.4013).

ACHTUNG: Wenn Sie Deutsch sprechen, stehen Ihnen kostenlos sprachliche Hilfsdienstleistungen zur Verfügung. Rufnummer: 1.800.940.5049 (TTY: 763.847.4013).

ប្រយ័ត្ន: បើសិនជាអ្នកនិយាយភាសាខ្មែរ, សេវាជំនួយភាសា ដោយមិនគិតថ្លៃ គឺអាចមានសំរាប់អ្នក។ ហៅ 1.800.940.5049 (TTY: 763.847.4013).

ملحوظة: إذا كنت تتحدث اذكر اللغة، فإن خدمات المساعدة اللغوية تتوافر لك بالمجان. اتصل برقم 1.800.940.5049 (رقم هاتف الصم والبكم: 763.847.4013).

ATTENTION : Si vous parlez français, des services d'aide linguistique vous sont proposés gratuitement. Appelez le 1.800.940.5049 (TTY: 763.847.4013).

주의: 한국어를 사용하시는 경우, 언어 지원 서비스를 무료로 이용하실 수 있습니다. 1.800.940.5049 (TTY: 763.847.4013). 번으로 전화해 주십시오.

PAUNAWA: Kung nagsasalita ka ng Tagalog, maaari kang gumamit ng mga serbisyo ng tulong sa wika nang walang bayad. Tumawag sa 1.800.940.5049 (TTY: 763.847.4013).